

y; meth2 arg2 X pabal y ad20; gal<sub>1</sub> sm phen2  
 ↓ bil (T1-free)  
 ↓ UV irradiation (5% survival)  
 rAl bil (T1-free)  
 ↓ UV irradiation (5% survival)  
 y; meth2 arg2 gal<sub>1</sub> sm phen2 X rAl bil; palB7 (T1-free)  
 ↓  
 bil; palB7

YOUR STOCK NUMBER FOR THIS CULTURE  
 ORIGIN OF STOCK:  
 ORIGINAL MUTANTS: Mutagen (UV (5% survival))  
 RECOMBINANTS: Cross of origin:

Pedigree of employed strains: Please use reverse side of this sheet for  
 data, or give reference to source of data.

Test for translocations: Tester strain:  
 translocation(s) present:  
 (designate by linkage group)  
 Tested by:

Published Reference(s):  
 DOW, G. (1963). Science 120: 1183-1184  
 (1963). Genet. Res. Camb. 6: 11-26

Please note unusual characteristics (i.e. genetic stability, growth and  
 scoring methods, etc.)  
 See Genet. Res. Camb. for method of scoring sporulation of most phenotypes.  
 Strains can be improved by elevating the level of sporangia production.

YOUR NAME (London form)  
 DATE SUBMITTED August 1, 1966  
 please do not write below this line

sent for:	Name	date	genotype:	viability:	lipidized:

Comments:

LINKAGE