

Kafw, Mardad + Cohen  
 Mutation freq: 38(19%) 141-146

TABLE 1: GENOTYPES OF STRAINS PRODUCING TEST DIPLOIDS

FGSC Numbers	Genotype	Diploid No
# 473	Su1A1 adE20; AcrA1 wA3; ActA1; pyroA4; facA303; lacA1 sB3; O11A2 choA1; riboB2 chaA1	2196
# 475	fpaB37 galD5 sua1adE20 riboA1 ana1 pabaA1 ya2 adE20 b1A1; sD85 fwa2	2196
# 513	Su1A1 adE20; AcrA1; ActA1; pyroA4; facA303; lacA1 sB3; choA1; chaA1	2100
<u>Strains for testing of sulfonilamides</u>		
# 515	fpaB37 galD5 sua1adE20 riboA1 ya2 adE20; pyroA4; facA303; chaA1	2199
# 514	ana1 <sup>+</sup> adE20 b1A1; AcrA1; ActA1; sB3; fwa2	2199

been made :  
 purgo's group  
 (a) In a  
 arm of  
 gregants is  
 which non-  
 ways of a li  
 diploid segre  
 several semi  
 zygous conc  
 ly recognize  
 effect can b  
 non-disjunct  
 few haploid  
 function [3  
 ter results n  
 (which may  
 ment with p  
 seo-fulvin [3  
 (b) If the  
 vegetatively  
 resulting fro  
 [17]. In all  
 are of fawn  
 tively, when  
 twin pairs a  
 tors (AcrA  
 2196), or y  
 over  
 sion  
 twin sector  
 and substan  
 identified v  
 or dispropo  
 facts [9,16  
 (c) To ch  
 diploids ca  
 letions are  
 growth, sp  
 can not on  
 recognized  
 tants of the  
 (d) If a  
 haploids ca  
 diploids wh  
 [15]. Unus  
 can be ide  
 markers be