Fungal Genetics Stock Center
Cell Biology and Biophysics
School of Biological Sciences
5007 Rockhill Road
University of Missouri, Kansas City
Kansas City, MO 64110

AndT

PLEASE PROVIDE COMPLETE INFORMATION

Reprints or other data relating to this deposit will aid the Stock Center and recipients of the strain.

Accession number

SPECIES Aspergillus nidulans	MATING TYPE N/A N/A
GENOTYPE pyrG89; argB2; pabaB22, nkuA::argB; riboB2	
DESIGNATION OF MUTANT ALLELE(S)	
LINKAGE GROUP(S) STRAIN DESIGNA	TION IF WILD-TYPE N/A
YOUR STOCK NUMBER FOR THIS CULTURE TNO include stock no. from other collecti	

ORIGIN OF STOCK The nkuA gene was deleted in KJ12 (wA3; argB2; pyroA4) by replacing it with argB. This was crossed to A770 (pabaB22; pyrG89; riboB2). TN02A25 is a segregant of this cross.

for example - obtained from, genetic background, from cross with; or if collected from nature, collection point, substrate and collector.

PUBLISHED REFERENCES Genetics (published ahead of print) available at http://www.genetics.org/aheadofprint.shtml

RECOMMENDED CATALOG LISTING Section C (Aspergillus)

IF UNPUBLISHED, please indicate strain of origin, mutagen, worker, genetic background, important characteristics N/A

COMMENTS (special growth requirements, aberrations, heterokaryon compatibility, special uses of strain, etc.)

Deletion of nkuA (the A. nidulans homolog of human Ku70 gene) reduces the frequency of non-homologous integration of transforming DNA and greatly facilitates gene targeting. NkuA deleted strains show no obvious growth defect and are not sensitive to a variety of DNA damaging drugs.

YOUR NAME TANIA NAYAK

DATE 01/03/2006