

plates of Vogel's medium N containing  
1.5% sorbose ; 0.2% glycerol and 0.2% glucose  
supplemented with 1mg/100mls of DL-  
p-fluorophenylalanine (FPA) strains which are  
not resistant to FPA (e.g. 74A) will  
not grow on these plates.

Fpr-1 is located between  
the isoleucine valine cluster and  
spray on IR. it is very tightly  
linked to act-2.

- T. Hayes (Texas A & M Univ) 11/1/75  
(ag) E. Koffi Bortny (Mc Gill U) 5/8/76  
(ag) M. A. Laper (U of Edinburgh, England) 12/30/76  
(ag) " " " " " " 4/11/77 (dupl)  
(ag) H. J. Darger (McMaster U) 8/9/77  
(ag) V. Prakash (U. of Malaya) 10/13/77  
(ag) F. Buxton (U of Leeds, Eng) 8/30/79  
(ag) M. Beal (Cornell U) 3/19/81  
(ag) J. C. Dunlop (U of Calif, Santa Cruz) 9/21/81  
(ag) M. Berrill (U of Sheffield) 8/30/82  
(ag) D. Perkins (Stanford U) 9/30/82  
(ag) I. Barthelme (U. Hannover) 11/1/83  
(ag) A. D. DeKink (Fla State U) 12/19/83