a component from 4-spored pseudohomothallic

Record of Neurospora Culture

Fungal Genetics Stock Center, Botany Department, Dartmouth College
Hanover, New Hampshire, U.S.A.

a
FGSC# 2511

GENOTYPE Hanalei-16 v 3 N. tetrasperma a
mating type

date 4/22/74

ALLELE DESIGNATION(S) ✓
(isolation no.)

YOUR STOCK NUMBER FOR THIS CULTURE P659

ORIGIN OF STOCK P556 A+a Collect by DDP from burnt grass, Kauai, June 1972
(for example - obtained from, induced in, from cross with, etc.)

~~Separated into~~ a component from conidial isolation by B. Turner.

PUBLISHED REFERENCE

(for data regarding origin, linkage, characteristics, etc.)

IF UNPUBLISHED, please indicate strain of origin, mutagen, worker, distinguishing characteristics

LINKAGE GROUP(S) ; COMMENTS (special growth conditions, aberrations, heterocaryon compatibility, genetic background, complementation group, etc.)

(use additional space on back of page if necessary)

YOUR NAME David D. Perkins DATE APR 19 74

Please do not write below this line

Please do not write in this space

opp mt = 2510

lyophilized 5/8/74

checked for viability 5/10/74 OC

checked for genotype

other storage method 5/3/74 grown onto complete

checked for viability 5/10/74 OC

sent to:

name	date	name	date
(sg) O. Mylyk (U of B.C., Canada)	7/23/76		
(sg) D. Natvig (U. New Mexico)	11/12/84		
(sg) J. Taylor (UC Berkeley	10/22/85		

Comments: fertile with black spores x tetrasperma A.
Obtained by conidial isolation from same culture (P556) a Hanalei-16v1
grows well on minimal