

Fungal Genetics Stock Center  
Dept. of Microbiology  
Univ. of Kansas Medical Center  
Kansas City, KS 66103

PLEASE PROVIDE COMPLETE INFORMATION

Reprints or other data relating to this deposit will aid the Stock Center and recipients of the strain.

Accession  
number

SPECIES N. crassa 7504

GENOTYPE bd cla-1

MATING TYPE a LINKAGE GROUP(S) IV; T.(I, VII)

DESIGNATION OF MUTANT ALLELE(S) \_\_\_\_\_

STRAIN DESIGNATION IF WILD-TYPE \_\_\_\_\_

YOUR STOCK NUMBER FOR THIS CULTURE 7107  
include stock no. from other collections

ORIGIN OF STOCK Cross #386: bd cla-1a (7060) x bd cla-1A (7061)  
for example - obtained from, genetic background, from cross with; or if  
collected from nature, collection point, substrate and collector.

PUBLISHED REFERENCES Brady et al (1988) Genome 30, 299  
(Abstract)

(for any information regarding this stock)

IF UNPUBLISHED, please indicate strain of origin, mutagen, worker,  
genetic background, important characteristics Originally phenotype seen in a nfa.  
Strain obtained from Perkins after irradiation of sn, cr

COMMENTS (special growth requirements, aberrations, heterokaryon  
compatibility, special uses of strain, etc.)

Translocation (I, VII) Breakpoint in I is between leu-3 + m.t.

(I) Prelim. mapping data leu-3 (120 m.u) cla-1 (13 m.u) m.t (111 m.u) csp-1  
(VII) " " shows cla-1 to be within 3-5 m.u. from oli<sup>2</sup>\*

(use additional space below or on back of page if necessary)

YOUR NAME S. BRODY DATE \_\_\_\_\_

Additional Comments: (use back of sheet if necessary)

\* The exact location in VII is unclear; ie which side of oli<sup>2</sup>.  
Keith Johnson (J. Dunlap's lab) had 4.5% recomb. between cla-1  
+ fig.