

Fungal Genetics Stock Center  
Dept. of Microbiology  
Univ. of Kansas Medical Center

Kansas City, KS 66103

PLEASE PROVIDE COMPLETE INFORMATION

Reprints or other data relating to this deposit will aid the Stock Center and recipients of the strain.

Accession  
number

SPECIES N.c. MATING TYPE A 8106

GENOTYPE ser-3

DESIGNATION OF MUTANT ALLELE(S) UK3-23

LINKAGE GROUP(S) IL

STRAIN DESIGNATION IF WILD-TYPE \_\_\_\_\_

YOUR STOCK NUMBER FOR THIS CULTURE 1837  
include stock no. from other collections

ORIGIN OF STOCK f<sub>1</sub> of original x I-marked tester. DP cross.

for example - obtained from, genetic background, from cross with; or if collected from nature, collection point, substrate and collector.

PUBLISHED REFERENCES \_\_\_\_\_

(for any information regarding this stock)

IF UNPUBLISHED, please indicate strain of origin, mutagen, worker,

genetic background, important characteristics Comes upon minimal after long lag.

50% white or pale ascospores x ser<sup>+</sup>. Detected as an ascospore color

mutant. Some Ser<sup>-</sup> spores darken with age.

COMMENTS (special growth requirements, aberrations, heterokaryon compatibility, special uses of strain, etc.)

Origin: In Am<sup>+</sup> transformant of am<sup>-</sup> ind A.

(See Perkins et al. 1993 Genetics 134:729-736).

Not a rearrangement.

(use back of page if necessary)

YOUR NAME David D. Perkins DATE 15 Dec 95